Subject Index for
Journal of Ichthyology, Volume 34, 1994

Abdominal wall: **3**, 144–149
Abiotic factors' effect: **7**, 104–118
Abramis brama: **7**, 62–69
Acanthopterygii: **9**, 27–72
Accidental introduction: **3**, 135–138
Acipenser baeri: **5**, 154–158
Acipenser baeri chatys: **8**, 58–71
Acipenser baeri stenorrhynchus: **8**, 58–71
Acipenser gueldenstaedti: **7**, 9–17
Acipenseridae: **2**, 90–103; **8**, 58–71; **9**, 111–124
Activity of hydrolases: **2**, 134–141
Adaptation to natural feeding: **1**, 96–106
Aerial olfaction: **6**, 38–47
Age: **4**, 67–73; **6**, 59–72
Age determination: **3**, 39–50
Alaskan pollack: **6**, 131–133; **7**, 92–103
Alepocephalidae: **2**, 104–111
Alepocephalus planifrons: **2**, 104–111
Aleyev, Yu.G.: **3**, 120–125
Anabantidae: **3**, 9–19
Anarhichas lupus: **1**, 140–151; **6**, 83–93
Ancient Lake El'gygytgyn: **2**, 20–34
Anguilliformes: **8**, 121–125
Antarctica: **4**, 67–73; **8**, 10–23
Antarctic cod: **4**, 67–73
Aphanopus intermedius larvae: **8**, 135–139
Arctic char: **3**, 90–103
Argentine squid fishery: **8**, 150–152
Astronesthidae: **7**, 142–150
Atherine fishes: **9**, 125–140
Atherinidae: **9**, 125–140
Atheriniform families: **9**, 27–72
Atherinomorpha: **9**, 27–72
Atlantic Ocean: **5**, 60–74
Atlantic salmon: **8**, 72–83
Autumn distribution: **6**, 48–58

Baikal whitefish: **2**, 69–77
Barbus: **7**, 119–141
Barents Sea: **4**, 112–127; **6**, 48–58
Bathygadidae: **5**, 98–115
Bathylagidae: **3**, 20–27
Baunt whitefish: **2**, 69–77
Behavioral mechanisms: **7**, 51–61
Behavior in unfamiliar environment: **3**, 139–143
Behavior reaction of amino acids: **2**, 90–103
Belligerent sculpin: **5**, 149–153
Bellingshausen Sea: **2**, 112–114
Beluga: **1**, 68–75
Beluga juveniles: **6**, 134–139
Bering Sea: **4**, 92–101; **6**, 73–82; **9**, 85–99
Bertelsen, Eric: **3**, 126–129
Bioenergetics: **6**, 83–93
Biological characteristics and distribution: **3**, 28–38
Biological status: **2**, 122–127
Biology of salmon shark: **2**, 115–121
Biotopic population distribution: **6**, 14–25
Black Sea: **6**, 149–151; **8**, 147–149; **9**, 125–140
Black Sea anchovy: **6**, 94–108
Black Sea basin: **6**, 14–25
Black Sea fishes: **8**, 35–43
Black Sea trout: **6**, 140–142
Blood plasma: **3**, 150–156
Blue whiting: **1**, 132–139
Bol'shaya River estuary: **1**, 76–83; **4**, 47–57; **5**, 149–153
Bony skeleton: **8**, 108–120
Brama australis: **5**, 144–148
Brama japonica: **1**, 107–116; **8**, 97–107
Bramidae: **8**, 97–107
Bream: **7**, 62–69
Breeding season: **9**, 73–84
Burbot breeding stocks: **7**, 18–28

Carcharhinus longimanus: **6**, 38–47
Careproctus dentatus: **2**, 8–19
Caspian Sea: **9**, 125–140
Caspian Sea basin: **6**, 14–25

Subject Index for
Journal of Ichthyology, Volume 34, 1994

Abdominal wall: **3**, 144–149
Abiotic factors' effect: **7**, 104–118
Abramis brama: **7**, 62–69
Acanthopterygii: **9**, 27–72
Accidental introduction: **3**, 135–138
Acipenser baeri: **5**, 154–158
Acipenser baeri chatys: **8**, 58–71
Acipenser baeri stenorrhynchus: **8**, 58–71
Acipenser gueldenstaedti: **7**, 9–17
Acipenseridae: **2**, 90–103; **8**, 58–71; **9**, 111–124
Activity of hydrolases: **2**, 134–141
Adaptation to natural feeding: **1**, 96–106
Aerial olfaction: **6**, 38–47
Age: **4**, 67–73; **6**, 59–72
Age determination: **3**, 39–50
Alaskan pollack: **6**, 131–133; **7**, 92–103
Alepocephalidae: **2**, 104–111
Alepocephalus planifrons: **2**, 104–111
Aleyev, Yu.G.: **3**, 120–125
Anabantidae: **3**, 9–19
Anarhichas lupus: **1**, 140–151; **6**, 83–93
Ancient Lake El'gygytgyn: **2**, 20–34
Anguilliformes: **8**, 121–125
Antarctica: **4**, 67–73; **8**, 10–23
Antarctic cod: **4**, 67–73
Aphanopus intermedius larvae: **8**, 135–139
Arctic char: **3**, 90–103
Argentine squid fishery: **8**, 150–152
Astronesthidae: **7**, 142–150
Atherine fishes: **9**, 125–140
Atherinidae: **9**, 125–140
Atheriniform families: **9**, 27–72
Atherinomorpha: **9**, 27–72
Atlantic Ocean: **5**, 60–74
Atlantic salmon: **8**, 72–83
Autumn distribution: **6**, 48–58

Baikal whitefish: **2**, 69–77
Barbus: **7**, 119–141
Barents Sea: **4**, 112–127; **6**, 48–58
Bathygadidae: **5**, 98–115
Bathylagidae: **3**, 20–27
Baunt whitefish: **2**, 69–77
Behavioral mechanisms: **7**, 51–61
Behavior in unfamiliar environment: **3**, 139–143
Behavior reaction of amino acids: **2**, 90–103
Belligerent sculpin: **5**, 149–153
Bellingshausen Sea: **2**, 112–114
Beluga: **1**, 68–75
Beluga juveniles: **6**, 134–139
Bering Sea: **4**, 92–101; **6**, 73–82; **9**, 85–99
Bertelsen, Eric: **3**, 126–129
Bioenergetics: **6**, 83–93
Biological characteristics and distribution: **3**, 28–38
Biological status: **2**, 122–127
Biology of salmon shark: **2**, 115–121
Biotopic population distribution: **6**, 14–25
Black Sea: **6**, 149–151; **8**, 147–149; **9**, 125–140
Black Sea anchovy: **6**, 94–108
Black Sea basin: **6**, 14–25
Black Sea fishes: **8**, 35–43
Black Sea trout: **6**, 140–142
Blood plasma: **3**, 150–156
Blue whiting: **1**, 132–139
Bol'shaya River estuary: **1**, 76–83; **4**, 47–57; **5**, 149–153
Bony skeleton: **8**, 108–120
Brama australis: **5**, 144–148
Brama japonica: **1**, 107–116; **8**, 97–107
Bramidae: **8**, 97–107
Bream: **7**, 62–69
Breeding season: **9**, 73–84
Burbot breeding stocks: **7**, 18–28

Carcharhinus longimanus: **6**, 38–47
Careproctus dentatus: **2**, 8–19
Caspian Sea: **9**, 125–140
Caspian Sea basin: **6**, 14–25

Catch coefficients: **3**, 61–72
Caucasian gobies: **6**, 14–25
Central Asia: **3**, 104–113
Central Chukotka: **2**, 20–34
Ceratoscopelus warmingii: **6**, 59–72
Channel catfish juveniles: **5**, 88–97
Char: **1**, 42–56; **2**, 20–34
Chernobyl: **4**, 16–38
Chronic intoxication: **6**, 134–139
Chum salmon: **4**, 92–101; **7**, 40–50
Cigarfishes: **7**, 70–82
Classification of chars: **4**, 128–147
Coastal waters: **5**, 12–26
Cod: **4**, 112–127; **6**, 83–93
Collections made under ice: **2**, 128–133
Commercial exploitation: **3**, 114–119
Comparative analysis: **9**, 111–124
Comparative characterization: **1**, 12–23
Comparative morphological analysis: **8**, 58–71
Comparative morphology: **2**, 35–47
Comparative restriction analysis: **2**, 69–77
Conceptual analysis: **7**, 83–91
Condition determination: **6**, 131–133
Congiopodidae: **2**, 149–155
Congridae: **3**, 73–89
Cooling pond: **4**, 16–38
Coregonid fishes: **6**, 117–130
Coregonus lavaretus baicalensis: **2**, 69–77
Coregonus lavaretus baunti: **2**, 69–77
Coregonus muksun: **7**, 104–118
Cottidae: **1**, 1–11; **4**, 1–15; **5**, 149–153
Cottocomephorus grewingki: **1**, 1–11; **1**, 12–23; **4**, 1–15
Craniological analysis: **2**, 35–47; **6**, 14–25
Crocodile shark: **4**, 155–157
Ctenopoma: **3**, 9–19
Cyclopteridae: **2**, 122–127
Cyprinidae: **3**, 104–113; **7**, 119–141; **8**, 24–34; **8**, 84–96
Cyprinus carpio: **3**, 144–149

Dace: **3**, 104–113
Deepest capture: **6**, 143–148
Dekhnik, Tat'yana Vladimirovna: **8**, 153–156
Density-dependent behavior: **9**, 73–84
Density regulation: **7**, 51–61
Derepodichthys alepidotus Gilbert: **8**, 126–131
Diet of major fishes: **8**, 150–152
Diet relationships: **6**, 149–151
Diet: **1**, 107–116; **4**, 92–101; **5**, 154–158; **6**, 73–82
Distribution: **3**, 73–89; **4**, 92–101; **5**, 149–153
Distribution characteristics: **2**, 122–127
Distribution of *Cubiceps*: **5**, 116–143
Distribution of Japanese mackerel: **4**, 74–91
Distribution of mesopelagic fishes: **7**, 1–8
Downstream transport of fish: **7**, 29–39

Early development stages: **7**, 62–69
Early stages of ontogeny: **5**, 60–74
Eastern Atlantic: **7**, 1–8
Ecology: **5**, 144–148
Eel genus: **3**, 73–89
Effect of season on age: **2**, 78–89
Effects of starvation: **1**, 96–106
Eleginopsidae: **8**, 10–23
Eleginus navaga: **6**, 48–58
Embryonic development: **5**, 27–45; **5**, 46–59
Embryonic-larval development: **1**, 42–56
Emergency sewage disposal: **7**, 18–28
Energy metabolism: **8**, 35–43
Engraulis encrasicolus ponticus: **6**, 94–108
Environmental factors of natural mortality: **6**, 109–116
Enzyme activity: **8**, 35–43
Equatorial Indian Ocean: **4**, 155–157
Erratum: **9**, 152–155
Error Seamount: **8**, 121–125
Ethiopia: **3**, 9–19
Ethiopian fauna: **7**, 119–141
Eumicrotremus soldatovi: **2**, 122–127
Exocoetidae: **5**, 12–26; **5**, 60–74
External structure of postorbital organ: **7**, 142–150

Factors determining spawner survival time, **1**, 57–67
Family Nototheniidae: **8**, 108–120

Family Rajidae: **6**, 26–37
Family Salmonidae: **2**, 20–34
Far Eastern smelt: **5**, 27–45
Far North seas: **6**, 26–37
Feeding: **1**, 76–83; **7**, 70–82
Feeding conditions: **1**, 117–131
Fertilization: **1**, 140–151
Field experiments: **9**, 73–84
Fish behavior: **9**, 1–26
Fish community: **4**, 47–57
Fish populations: **3**, 114–119
Fish stocking density: **4**, 39–46
Fishes of northern latitudes: **2**, 134–141
Fluctuating asymmetry: **4**, 148–155
Flying fish: **5**, 12–26
Food search: **6**, 38–47
Forgotten species *Amitra liparina*: **2**, 1–7
Formation of morphological differences: **7**, 119–141
Fossil Notothenioid: **8**, 10–23
Fractional composition: **9**, 111–124
Freshwater ichthyofauna: **1**, 24–41

Gadiform: **8**, 10–23
Gadiformes: **5**, 98–115
Gadus morhua morhua: **4**, 112–127; **6**, 83–93
Gavialiceps: **3**, 73–89
Gear for sampling young fishes: **3**, 61–72
Genetic monitoring: **7**, 40–50
Genetic relationships: **6**, 117–130
Genetic similarity: **2**, 60–68
Genus *Astronesthes*: **7**, 142–150
Genus *Atherina*: **9**, 125–140
Genus *Cubiceps*: **7**, 70–82
Genus *Salvelinus*: **2**, 48–59
Gigantactinidae: **8**, 132–134
Gigantactis elsmani: **8**, 132–134
Gobiidae: **6**, 14–25
Goby subgenus: **2**, 35–47
Gonadotropic hormones: **3**, 150–156
Gray mullet: **3**, 150–156
Growth: **4**, 67–73; **6**, 59–72
Growth rate: **8**, 97–107
Growth-rate estimate: **3**, 39-50
Growth stimulation: **2**, 142-148
Gulf of California: **8**, 126–131
Guppies: **3**, 139–143
Gvozdarus sevtovidovi gen. et sp. n.: **1**, 152–158

Hatchery-raised chinook salmon: **1**, 96–106
Hatchery-raised chum salmon: **1**, 96–106
Hirundichthys affinis: **5**, 60–74
Hirundichthys speculiger: **5**, 60–74
Historical perspectives: **9**, 27–72
Huso huso: **1**, 68–75
Hydrothermal zone inhabitant: **8**, 126–131
Hypomesus olidus: **5**, 27–45
Hypophthalmichthys: **5**, 75–87
Hypophthalmichthys molitrix: **1**, 84–95; **4**, 16–38
Ichthyofauna: **9**, 141–151
Ichthyophthiriasis: **5**, 88–97
Ictalurus punctatus: **5**, 88–97

Indian Ocean: **6**, 59–72; **8**, 1–9
Indian Ocean sector: **4**, 67–73
Indigirka River: **5**, 154–158
Individual development rates: **8**, 108–120
Individual electronic activity: **8**, 147–149
Indo-West Pacific: **3**, 73–89
Industrial races: **8**, 84–96
Intensive rearing: **5**, 88–97
Interspecific comparisons: **6**, 152–156
Intraluminal and membrane digestive processes: **2**, 134–141
Intraspecific comparisons: **6**, 152–156
Intraspecific differentiation: **1**, 12–23
Irkutsk Reservoir: **3**, 135–138

Japanese mackerel: **4**, 74–91
Juvenile fishes: **6**, 149–151
Juvenile sockeye salmon: **4**, 102–111
Juvenile sturgeons: **2**, 90–103

Kamchatka: **4**, 92–101; **4**, 102–111
Kamchatkan Pacific waters: **7**, 92–103
Kamchatkan trout: **2**, 60–68
Karyological analysis: **6**, 14–25
Karyotypic diversity: **6**, 1–13
Kazakhstan: **3**, 104–113

Kolyma River: **5**, 154–158
Krefft, Gerhard: **3**, 130–134

Lake Achchen: **5**, 1–11
Lake Azabach'e: **4**, 58–66; **4**, 102–111
Lake El'gygytgyn: **2**, 20–34; **4**, 128–147
Lake Ladoga: **1**, 42–56
Lake Onega: **1**, 42–56
Lakes of Chukotsk Peninsula: **2**, 48–59
Lamna ditropis: **2**, 115–121
Large African barbs: **7**, 119–141
Larval and juvenile fishes: **2**, 128–133
Larval capelin: **4**, 112–127
Larval carp: **3**, 144–149
Larval development: **5**, 27–45
Larval fishes: **6**, 149–151
Late Eocene Seymour Island: **8**, 10–23
Learning in fish: **6**, 152–156
Lena River: **8**, 58–71
Length-age structure: **8**, 97-107
Length of freshwater residence: **4**, 58-66
Leuciscinae: **8**, 24–34
Leuciscus Agassiz: **3**, 104–113
Leuciscus aphipsi Aleksandrov: **8**, 24–34
Leuroglossus schmidti: **3**, 20–27
Liopsetta glacialis: **3**, 1–8
Liparidae: **2**, 1–7; **2**, 8–19; **8**, 1–9
Liver: **6**, 134–139
Liza ramada: **3**, 150–156
Lophiformes: **8**, 132–134
Lota lota: **7**, 18–28
Lower Volga: **1**, 68–75

Macrouridae: **5**, 98–115
Macrourids: **5**, 98–115
Mallotus villosus villosus: **4**, 112–127
Marine organisms: **8**, 44–57
Matrices for estimating stocks: **8**, 44–57
Matrices for predicting catch: **8**, 44–57
Maturation: **5**, 75–87
Maturation of yearlings: **6**, 94–108
May spawning stock: **4**, 1–15
Megalocottus platycephalus platycephalus: **5**, 149–153
Mesopelagic fishes: **6**, 73–82; **7**, 1–8; **9**, 85–99
Micromesistius poutassou: **1**, 132–139
Migration of Japanese mackerel: **4**, 74–91
Mitochondrial DNA: **2**, 69–77
Morphobiological forms: **7**, 104–118
Morphoecological differentiation: **8**, 72–83
Morphofunctional condition: **6**, 134–139
Morphological characteristics: **3**, 9–19; **3**, 144–149
Morphological differentiation: **7**, 92–103
Morphological divergence: **8**, 84–96
Moscow: **8**, 84–96
Moscow city boundaries: **9**, 141–151
Moscow River: **9**, 141–151
Motor activity in fish: **6**, 152–156
Mozambique: **2**, 142–148
Muksun: **7**, 104–118
Muscle tissue: **6**, 134–139
Myctophidae: **6**, 59–72; **7**, 151–160
Myctophids distribution: **7**, 151–160

Natural mortality estimation: **6**, 109-116
Natural production effectiveness: **7**, 9-17
Natural reproduction: **1**, 68-75
Navaga: **6**, 48–58
Neogobius kessleri: **2**, 35–47
Neogobius ratan: **2**, 35–47
Neogobius syrman: **2**, 35–47
Nettastomatidae: **8**, 121–125
Nettenchelys erroriensis sp. nov.: **8**, 121–125
New genus: **2**, 8–19
New species: **2**, 104–111
New species of *Paraliparis*: **8**, 1–9
New species *Paraliparis challengeri*: **2**, 1–7
Nine-spined stickleback: **5**, 46–59
Nomeidae: **5**, 116–143
North American trout: **2**, 60–68
North Pacific: **8**, 135–139
Northeast Atlantic: **2**, 1–7
Northeastern Atlantic: **1**, 132–139
Northeastern Sea of Okhotsk: **2**, 122–127
Northern Karelia: **8**, 72–83
Northwest Pacific Ridge: **5**, 98–115
Northwestern Indian Ocean: **8**, 121–125
Northwestern Pacific: **3**, 20–27

Northwestern Pacific Ocean: **2**, 115–121; **4**, 74–91
Norwegian Sea: **4**, 112–127; **6**, 143–148
Notalepis annulata: **2**, 112–114
Notothenia rossii rossii: **4**, 67–73
Nototheniidae: **1**, 152–158
Notothenioidei: **8**, 10–23
Nuclear power plant: **4**, 16–38

Ob Bank: **4**, 67–73
Ob River: **3**, 51–60
Oceanic whitetip shark: **6**, 38–47
Oncorhynchus: **1**, 96–106; **7**, 51–61; **9**, 73–84
Oncorhynchus gorbuscha: **7**, 40–50; **7**, 83–91; **9**, 73–84
Oncorhynchus keta: **4**, 92–101; **7**, 40–50
Oncorhynchus mykiss: **2**, 78–89; **3**, 135–138; **8**, 140–146
Oncorhynchus nerka: **1**, 57–67; **1**, 117–131; **4**, 58–66; **4**, 102–111; **7**, 40–50; **7**, 51–61
Oncorhynchus tshawytscha: **1**, 96–106
Ontogenesis: **4**, 1–15
Oreochromis mossambicus: **2**, 142–148
Osmeridae: **5**, 27–45

Pacific Ocean: **4**, 92–101
Pacific pomfret: **1**, 107–116; **8**, 97–107
Pacific salmon: **4**, 148–155; **7**, 51–61; **9**, 73–84
Pakhacha River populations: **7**, 40–50
Paralepididae: **2**, 112–114
Patagonian shelf: **8**, 150–152
Perciformes: **6**, 1–13; **8**, 10–23
Peruvian jack mackerel: **3**, 39–50
Phylogenetic relationships: **4**, 128–147
Pink salmon: **7**, 40–50; **7**, 83–91; **9**, 73–84
Pisces: **1**, 152–158
Plaice: **6**, 83–93
Platichthys stellatus: **1**, 76–83
Pleuronectes platessa: **6**, 83–93
Poecilia reticulata: **3**, 139–143
Polar dab: **3**, 1–8
Pond raising techniques: **3**, 144–149
Ponticola Iljin: **2**, 35–47
Population: **5**, 149–153
Population organization: **7**, 83–91
Populations of small water bodies: **8**, 84–96
Porcupine Seabight: **2**, 1–7
Post-disaster period: **4**, 16–38
Postorbitals: **4**, 148–155
Previtellogenous oocyte ultrastructure: **8**, 140–146
Production: **6**, 59–72
Production of Siberian roach: **3**, 51–60
Proeleginops grandeastmanorum gen. et sp. nov.: **8**, 10–23
Pseudocarcharias kamoharai: **4**, 155–157
Pseudocarchariidae: **4**, 155–157
Pungitius pungitius: **5**, 46–59

Quadruped locomotion: **2**, 149–155
Qualitative changes: **2**, 78–89
Quantitative changes: **2**, 78–89

Radiation suppression of gonads in females: **2**, 142–148
Radioactive contamination: **4**, 16–38
Radiobiological analysis: **4**, 16–38
Rainbow trout: **3**, 135–138
Raja clavata: **8**, 147–149
Raja radiata: **6**, 143–148
Rapid method: **6**, 131–133
Redescription: **8**, 24–34
Regulation of Volga: **7**, 9–17
Reproduction: **1**, 42–56
Reproductive biology: **1**, 1–11; **6**, 26–37
Reproductive characteristics: **1**, 132–139
Reproductive cycle: **4**, 1–15
Reproductive potential of fishes: **9**, 100–110
Reproductive system of fish: **4**, 16–38
Ringed Notolepis: **2**, 112–114
Roach: **8**, 84–96
Role of heterochrony: **7**, 119–141
Ross Sea: **1**, 152–158
Russian study: **9**, 1–26
Russian sturgeon: **7**, 9–17
Russian sturgeon juveniles: **6**, 134–139
Rutilus rutilus: **8**, 84–96
Rutilus rutilus caspicus: **3**, 28–38
Rutilus rutilus lacustris: **3**, 51–60
Rybinsk Reservoir: **9**, 100–110

Rybinskoe Reservoir: **7**, 18–28

Salmo salar: **8**, 72–83
Salmo trutta: **8**, 72–83
Salmo trutta labrax: **6**, 140–142
Salmon shark: **2**, 115–121
Salvelinus: **3**, 90–103
Salvelinus alpinus lepechini: **1**, 42–56
Sarcoplasmic muscle proteins: **9**, 111–124
Scale growth annual variation: **1**, 117–131
Scomber japonicus: **4**, 74–91
Scorpaeniformes: **1**, 1–11; **2**, 1–7; **2**, 8–19; **2**, 149–155; **8**, 1–9
Sea of Azov: **6**, 140–142; **9**, 125–140
Sea of Okhotsk: **2**, 8–19; **8**, 132–134
Sea trout: **8**, 72–83
Seasonal distribution: **2**, 115–121
Selection: **7**, 62–69
Sensory systems: **9**, 1–26
Sexual cycle: **1**, 12–23; **3**, 150–156
Sheksna reach: **9**, 100–110
Sheksninskii reach: **7**, 18–28
Siberia: **6**, 117–130
Siberian roach: **3**, 51–60
Siberian sturgeon: **5**, 154–158
Siberian sturgeon subspecies: **8**, 58–71
Silver carp: **1**, 84–95; **4**, 16–38; **5**, 75–87
Size–weight relationships: **6**, 149–151
Skate: **8**, 147–149
Skate groups: **8**, 147–149
Skates: **6**, 26–37
Skull development during ontogeny: **3**, 1–8
Skull morphology: **9**, 125–140
Slickhead: **2**, 104–111
Small African barbs: **7**, 119–141
Small streams: **8**, 72–83
Smoothtongue: **3**, 20–27
Snailfishes: **2**, 8–19
Sockeye salmon: **1**, 57–67; **1**, 117–131; **4**, 58–66; **7**, 40–50; **7**, 51–61
Southeastern Caspian Sea: **3**, 28–38
Southeastern India: **5**, 12–26
Southeastern Pacific: **2**, 104–111; **5**, 144–148
Southern pomfret: **5**, 144–148
Southern Sakhalin: **1**, 24–41
Southwest Atlantic: **7**, 151–160
Spawner survival time: **1**, 57–67
Spawners: **1**, 1–11; **9**, 73–84
Spawning: **5**, 75–87
Spawning of yearlings: **6**, 94–108
Spawning stocks: **1**, 12–23
Species-specific metabolites: **4**, 39–46
Sperm production: **2**, 78–89
Squaloliparis: **2**, 8–19
Stability of biological systems: **3**, 114–119
Starry flounder: **1**, 76–83
Steelhead: **2**, 78–89
Steroidal hormones: **3**, 150–156
Stocks of fishes: **9**, 100–110
Stress prevention: **1**, 84–95
Stress treatment: **1**, 84–95
Stromateoidei: **7**, 70–82
Sturgeon representatives: **9**, 111–124
Subantarctic islands: **8**, 1–9
Suborder Notothenioidei: **6**, 1–13
Summer daily feeding dynamics: **9**, 85–99
Survival strategies: **3**, 90–103
Sympatric char: **2**, 48–59
Synthetic gonadotropin-releasing hormone: **5**, 75–87
Systematic relationships: **9**, 27–72
Systematics: **3**, 73–89; **3**, 104–113; **6**, 14–25
Systematics of *Cubiceps*: **5**, 116–143
Systematics problems: **9**, 125–140

Taranets char: **5**, 1–11
Taxonomic diversity: **6**, 1–13
Temperature conditions: **1**, 117–131
Temperatures in annual cycle: **8**, 35–43
Theragra chalcogramma: **6**, 131–133; **7**, 92–103
Thorny skate: **6**, 143–148
Tilapia: **2**, 142–148
Trachurus symmetricus murphyi: **3**, 39–50
Transliteration of Cyrillic alphabet: **6**, i
Tribe Salvelinini: **2**, 20–34
Trichiuridae: **8**, 135–139
Trophic differentiation: **4**, 102–111
Tropical zone: **6**, 59–72

Ust'-Khantaiskoe Reservoir: **7**, 29–39

Variability of biological indicators: **1**, 1–11
Volga–Caspian district: **7**, 62–69

Water dynamics: **7**, 151–160
Water structure: **7**, 151–160
Weddell Sea: **2**, 128–133
West Bering Sea: **7**, 92–103
Western Kamchatka: **1**, 76–83; **4**, 47–57; **5**, 149–153
White Nile basin: **3**, 9–19
Wolffish: **1**, 140–151; **6**, 83–93
World Ocean: **5**, 116–143

Yellowfin Baikal sculpin: **1**, 1–11; **1**, 12–23; **4**, 1–15
Yenisey River: **8**, 58–71
Young-of-the-year vobla: **3**, 28–38

Zanclorhynchus spinifer: **2**, 149–155
Zoarcidae: **8**, 126–131